





Product Information Sheet

Biatain Ibu Non-Adhesive Foam / Biatain Ibu Soft-Hold Foam	
Classification	Pain Control Dressing: Ibuprofen
British Columbia Practice	<ul style="list-style-type: none"> Consult with NSWOC / Wound Clinician / Physician / NP prior to use.
Key Points	<ul style="list-style-type: none"> Polyurethane foam dressing containing ibuprofen (0.5 mg/cm²) dispersed throughout dressing. Soft, conformable and highly absorbent. Provides an optimal moist wound healing environment and effective exudate management. Ibuprofen is released into the wound bed when in contact with wound exudate. Semi-permeable film backing is waterproof and provides a bacterial barrier. Two versions available: <ol style="list-style-type: none"> Biatain Ibu Non-Adhesive Foam Dressing is suitable for use on fragile skin due to absence of adhesive. Biatain Ibu Soft-Hold Foam Dressing has a partially adherent layer enabling dressing to stay in place while secondary dressing or compression therapy is applied. Must be in contact with wound bed and will conform to wound depth of up to 2 cm. Primary dressing but does not require a secondary dressing.
Indications	<ul style="list-style-type: none"> Painful wound. Biatain Ibu Non-Adhesive Foam dressing can be used on donor sites. Can be used in combination with compression therapy.
Precautions	<ul style="list-style-type: none"> Clinically infected wounds must be treated prior to use of Biatain Ibu Foam. Do not exceed 1200 cm² (e.g. 3 dressings of 20 x 20 cm each) at each dressing change. Dressing should not be changed more than twice daily corresponding to maximum daily use of 2400 cm². Can be used for up to 6 weeks as long as clinically indicated. Remove dressing prior to radiation treatment or examinations (including, x-rays, ultrasonic treatment, diathermy, microwaves or MR scanning). Has not been evaluated on lactating individuals, consult with physician / NP prior to using on this population.
Contraindications	<ul style="list-style-type: none"> Sensitivity or allergy to ibuprofen, any other components of the dressing, acetylsalicylic acid or other NSAIDs. Untreated clinically infected wounds. Do not use Biatain Ibu Soft-Hold Foam dressings on donor sites. Do not use with oxidizing solutions (e.g. hypochlorite and hydrogen peroxide solutions). Pregnant individuals. Children under 12 years of age except under the direction of a physician / NP.
Formats & Sizes	<ul style="list-style-type: none"> Biatain Ibu Non-Adhesive Foam <ul style="list-style-type: none"> 10 x 10 cm 10 x 20 cm 15 x 15 cm 20 x 20 cm Biatain Ibu Soft Hold Foam <ul style="list-style-type: none"> 10 x 10 cm 20 x 20 cm <div style="text-align: center;">  </div>

Directions	Rationale / Key Points
Selection	
Select dressing size where foam pad overlaps wound edge by minimum of 2 cm. Dressing can be cut to size if needed.	For small sized dressings an overlap of 1 cm may be necessary. Incorrect sizing of dressing will adversely affect dressing function.
Preparation	
Cleanse wound and periwound / surrounding skin with sterile	The safety of other cleansing agents in combination with



Product Information Sheet

Directions	Rationale / Key Points
<p>normal saline or water.</p> <p>Dry periwound / surrounding skin.</p> <p>If required, apply barrier film to periwound skin.</p>	<p>Biatain Ibu Foam has not been demonstrated. See Wound Cleansing Procedure or QR Code below.</p> <p>Ensure that solution has dried prior to dressing application to minimize risk of periwound maceration.</p> <p>To protect periwound skin from moisture associated skin damage and medical adhesive related skin injury.</p>
Application	
<p>If there is minimal exudate, moisten Biatain Ibu Foam with sterile saline or apply amorphous wound gel to wound bed.</p> <p>For wounds with depth 2 cm or less place dressing directly over the wound with the plain, non-printed side toward the wound.</p> <p>Secure with tape or wrap with kling.</p>	<p>Ibuprofen is released to wound bed when in contact with wound exudate.</p> <p>Must be in contact with wound bed and will conform to wound depth of up to 2 cm.</p> <p>The outer side prevents strike through of exudate.</p> <p>The non-printed side will release Ibuprofen into wound.</p>
Removal	
<p>Consider using adhesive remover to remove adhesives (e.g. tape).</p> <p>Gently lift corners of dressing away from wound.</p>	<p>To decrease risk of medical adhesive related skin injury (MARSI).</p> <p>To avoid trauma to wound and periwound skin</p>
Frequency of Dressing Change	
<p>May be left in place for up to 7 days if pain is effectively managed. Usual wear-time is 2- 3 days.</p> <p>Change dressing when clinically indicated or when visible signs of exudate approach edge of dressing.</p>	<p>Dressing change frequency is dependent on amount of wound exudate.</p> <p>Absorbed exudate is clearly visible through backing of dressing.</p>
Expected Outcomes	
<p>Wound pain is managed for 2-3 days</p> <p>Product performs as expected.</p>	<p>If product does not perform as expected notify NSWOC/Wound Clinician and consider submitting a Supply Chain Product Concern form.</p>
QR Codes	
 <p>Wound Cleansing</p>	
For further information please contact NSWOC/Wound Clinician	