




Product Information Sheet

| Remedy Prevent | |
|----------------------------|---|
| Classification | Skin Care: Protectant - Silicone |
| Key Points | <ul style="list-style-type: none"> • Remedy Clinical Prevent Silicone Cream is made with 24% silicone blended with botanical nutrients to provide moisturizing protective barrier. • May be used on intact and non-intact skin. • Fragrance-free, hypoallergenic, paraben-free, sulfate-free. • Can be used in the presence of Chlorhexidine Gluconate (CHG). • Does not interfere with the integrity of latex, nitrile, and vinyl gloves. |
| Indications | <ul style="list-style-type: none"> • For all ages of patients/clients/residents for the: <ul style="list-style-type: none"> ○ Prevention and treatment of mild/moderate Incontinence Associated Dermatitis (IAD). ○ Prevention and treatment of Moisture Associated Skin Damage (MASD) around wounds, tubes/ drains. ○ Treatment of chafed, chapped or cracked skin. |
| Precautions | <ul style="list-style-type: none"> • When cream is used on the feet: <ul style="list-style-type: none"> ○ Do not apply cream between the toes; remove any cream that may inadvertently get between toes. ○ Ensure socks and shoes are worn when walking to prevent a fall. |
| Contraindications | <ul style="list-style-type: none"> • Do not use for clients with known sensitivity or allergy to product ingredients • Do not use on wounds e.g. Stage 2 Pressure Injury • Do not use on severe Incontinence-Associated Dermatitis (IAD) • Do not use with silicone-based dressings or tapes |
| Formats & Sizes | <ul style="list-style-type: none"> • Tube <ul style="list-style-type: none"> ○ 59mL ○ 256mL <div style="text-align: center;">  </div> |

| Directions | Rationale / Key Points |
|---|---|
| Selection | |
| Select the size of tube appropriate to the client situation, (e.g., short stay consider the 59mL sized tube). Label the tube with date and the client's name. | The tube is for single-patient use only and is to be used within three years of opening. |
| Application | |
| Apply a thin layer to clean skin and gently smooth in until cream is absorbed. Prevention of IAD/Treatment of mild/moderate IAD: Apply a thin layer to skin that may come in contact with urine, stool or moisture/exudate. Treatment of dry, chafed, chapped, cracked skin: <ul style="list-style-type: none"> • Apply a thin layer to dry, chafed, chapped or cracked skin. • For feet, do not apply cream between the toes; remove any cream that may inadvertently get between toes. Prevention/Treatment of Periwound, Peri-tube/drain MASD: Allow cream to dry for 1-2 minutes before apply a dressing. Note: cannot be used with silicone-based dressings or tapes. | <p>Should not be applied so thick that there is a white layer and that the skin is not visible. A thin, transparent application provides good protection and allows for visual inspection of the skin.</p> <p>Product may make the soles of the feet slippery; ensure that socks/shoes are worn to prevent a fall.</p> <p>The silicone in the barrier cream may prevent the silicone cover dressings/tapes from adhering properly. This effect may also occur with other types of cover dressings/tapes</p> |



Product Information Sheet

| Directions | Rationale / Key Points |
|--|--|
| <p>Frequency of Application</p> | |
| <p>After each incontinent episode, apply to clean and dry skin.</p> <p>Dry, chafed, chapped or cracked skin; reapply at least twice/day.</p> | |
| <p>Expected Outcomes</p> | |
| <p>Prevention of IAD.</p> <p>Resolution of mild /moderate IAD. If no resolution of IAD within 3 days, or condition worsen, then consider the use of a zinc-based cream and other continence strategies.</p> <p>Resolution of dry, chafed, chapped, cracked skin.</p> <p>Prevention/resolution of periwound, peri-tube/drain MASD.</p> <p>Product performs as expected.</p> | <p>If product does not perform as expected, notify NSWOC/Wound Clinician and then consider submitting a Supply Chain Product Concern Form.</p> |
| <p>For further information please contact NSWOC/Wound Clinician</p> | |